

## Determination of zinc, copper, lead and cadmium concentration in breastmilk by anodic stripping voltammetry method and investigating their impact on infants' growth indicators

Naficeh Sadeghi<sup>1,2\*</sup>, Behrooz Jannat<sup>2</sup>, Masoomeh Behzad<sup>1</sup>, Mohammad Reza Oveisi<sup>1</sup>,  
Mannan Hajimahmoodi<sup>1</sup>, Fatemeh Ahmadi<sup>1</sup>

1- Department of Drug and Food Control, School of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

2- Halal Research Center, Ministry of Health and Medical Education, Tehran, Iran.

This paper is open access under [Creative Commons Attribution-NonCommercial 4.0 International](https://creativecommons.org/licenses/by-nc/4.0/) license.



**Submission:** 13 April 2020

**Revision:** 18 August 2020

**Acceptance:** 31 October 2020

### Abstract

**Background and objective:** Breastmilk provides nutritional, immunological, psychological, and developmental benefits for infants. Zinc and copper are of essential minerals, being involved in various biological processes. Their insufficient intake by infants can detrimentally affect function of body. In comparison, lead and cadmium are harmful trace elements which accumulate in women before pregnancy, transfer to fetus through placenta in pregnancy and then to infant through breastmilk in lactating period. The objective of our work was to investigate impact of zinc, copper, lead, and cadmium in breastmilk on infants' growth and their correlations with the mothers' dietary intake.

**Materials and methods:** Breastmilk samples and food intake information were achieved from 160 lactating women in Tehran. Concentration of copper, zinc, lead and cadmium were determined by anodic stripping voltammetry conducted by microwave digestion.

**Results and conclusion:** According to the results, concentration of copper, zinc, lead and cadmium were calculated as  $0.36 \pm 0.33$  mg/l,  $2.40 \pm 2.02$  mg/l,  $7.15 \pm 5.96$   $\mu$ g/l, and  $1.07 \pm 1.14$   $\mu$ g/l, respectively. Amounts of zinc and copper in breastmilk were increased by consumption of dairy, vegetables, fruits, bread, and nuts. Infants' height, weight and head circumference at birthday were directly correlated with zinc concentration while reversed correlation was observed for copper, lead and cadmium in the samples. In conclusion, controlled dietary intake of women before pregnancy and after childbirth has significant impact on healthy status of the infants. Obviously, more attention should be paid on potential sources of lead intake in women of Tehran because of its high concentration in the samples.

**Keywords:** Cadmium, copper, human breastmilk, lead, voltammetry, zinc

### 1. Introduction

Quality of the foods taken at early ages determines health status of human in adulthood.

Therefore, both nutrients and chemical contamination of human breastmilk is of concern for child healthcare practitioners. As reported,

\* Correspondence to: Naficeh Sadeghi; e-mail: [nsadeghi@sina.tums.ac.ir](mailto:nsadeghi@sina.tums.ac.ir); Tel.: +98-21-66954714, Fax: +98-21-66954714

amount of maternal zinc retained by fetus during the last trimester of pregnancy is less than the amounts secreted into the mothers' milk at the first three months of lactation [1]. Scientific evidences indicate that benefit of breastfeeding in favor of neural development attenuate the potential adverse impact of environmental neurotoxicants, through immunological, physiological, nutritional, and psychological advantages provided by breastmilk [2]. World Health Organization recommends just breastfeeding for nourishment of infants within the first six months of age followed by breastfeeding with complementary foods within two years of age [3].

Composition of human breastmilk is important for infants' development. Deficiency of essential trace elements results in impaired growth and development during infancy. Zinc deficiency can result in stunted growth and compromise immune function. Copper exists in all tissues and involves in red blood cells synthesis, maintenance of nerve cells, and function of immune system. It is recommended to intake 8-11 mg per day zinc and 0.9 mg per day copper in adulthood; even though they are recommended as 11 and 12 mg per day zinc and also 1 and 1.3 mg per day copper in pregnancy and lactation, respectively [4]. In comparison, contamination of breastmilk with toxic chemicals is concerned. Cadmium as a heavy metal widely disperses in the environment. Women are exposed to cadmium primarily by smoking or eating the foods contained/contaminated by it [5]. In addition, lead is another potentially toxic heavy metal with hepatotoxic, neurotoxic and nephrotoxic effects. It can also cause hemolytic anemia by disruption of cell membrane. Pregnant women and children who live near polluted regions (containing cathode ray tube (CRT) televisions and monitors, desktops, laptops, liquid crystal display (LCD) monitors, cell phones, keyboards, computer mice, printers, and copiers) are at severe risk of exposure to heavy metals such as cadmium and lead [6]. World Health Organization set a limit of 5 µg/l

for lead and 0.1 µg/l for cadmium in breastmilk [7].

In lactation, heavy metals are transported from the mothers' plasma to mammary glands and then secreted into breastmilk. In this regard, Kippler et al. reported that low exposure to cadmium by breastmilk at early ages induces oxidative stress [8]. Despite the study shown no impact of maternal diet on concentration of iron, zinc, and copper in the breastmilk [5], Honda et al. reported that amount of toxic metals in breastmilk collected from 68 Japanese mothers, aged 19-38 years, at 5-8 days postpartum was significant [9]. Therefore, impact of the mothers' nourishment on quality and quantity of their milk is a controversial issue [10].

Several techniques have been used for analysis of trace metals including flame atomic absorption spectrometry and inductively coupled plasma mass spectrometry (ICP-MS) [11,12]. Among them, differential pulse anodic stripping voltammetry (DPASV) has demonstrated great sensitivity for detection of zinc, copper, lead, and cadmium at low quantities. Therefore, this study aimed to determine amounts of zinc, copper, lead and cadmium by DPASV technique in breastmilk samples collected from 160 women within five months postpartum in Tehran in 2015. In addition, correlation of the heavy metals with infants' growth and dietary intake of the mothers is discussed.

## 2. Materials and methods

### 2.1. Sample size and sampling procedure

In the current research, the women who took nutritional supplements were excluded. Breastmilk of 160 healthy lactating women of Fazel, Kan, Yaftabad, and Arash Clinic in Tehran (Iran) were collected. About 10 ml of each sample was stored in acid-washed tube and kept at -20 °C until analysis.

The sample size (n) was calculated by Eq. 1.

$$n = \sigma^2(z_{1-\beta} + z_{1-\alpha})^2 / (\mu_0 - \mu_1)^2 \quad \text{Eq. 1}$$

$$\sigma = 2.02 \quad \alpha = 0.05 \quad \beta = 0.2$$

$$\mu_0 = 2.79 \quad \mu_1 = 2.40$$

## 2.2. Sample preparation

For mineralization, the samples were transferred to polyethylene tubes and digested by high performance microwave (Berghof Speed wave, Germany) equipped with a rotor designed for pressures up to 50 bar. Then, one ml of breastmilk was transferred to the digestion vessel and mixed with 5 ml nitric acid (65%, Suprapur®, Merck). Then, 2 ml of high purity hydrogen peroxide (30%, Suprapur®, Merck) was added and mineralization was done at 175 °C for 50 min. Hydrogen peroxide was further removed by heating and the solution was diluted by 5 ml of 0.1 M nitric acid.

## 2.3. Instrumental analysis

Amounts of zinc, copper, lead and cadmium were determined by differential pulse anodic stripping voltammetry with a hanging mercury drop electrode (Model 746 VA Trace Analyzer, Metrohm, Switzerland). The analysis was conducted by deposition potential of -1.20 V, deposition time of 30-90 s, scan rate of 6 mv/s, pulse amplitude of 50 mv, and final potential of +0.3 v. Standard solutions of zinc, copper, lead and cadmium at concentration of 1 g/l were prepared. At first, blank polarogram of the supporting electrolyte was recorded by using 10 ml of acetate buffer. Then, 500 µl of the samples were added to the cells and their polarogram was recorded. The analysis was done by standard addition method by adding 100 µl of the standard solutions to the samples [13].

## 2.4. Statistical analysis

The data was analyzed by SPSS software ver. 16. One-way ANOVA method was performed and the means were compared by Tukey test. All of the experiments were repeated three times. Difference of data was significant at  $p < 0.05$ . The descriptive data is presented as mean  $\pm$ SD.

## 3. Results and discussion

In the present study, limits of detection of 1.20 µg/l for zinc, 2.80 µg/l for copper, 0.30 µg/l for lead and 0.15 µg/l for cadmium were achieved. Results of the heavy metals are shown in Table 1, and descriptive data of the lactating women and their infants is presented in Table 2.

Table 1- Amounts of heavy metals in breastmilk of lactating women in Tehran

Clinic	N	Zn (mg/l)	Cu (mg/l)	Pb (µg/l)	Cd (µg/l)
A	40	1.33 $\pm 1.35$	0.64 $\pm 0.77$	6.32 $\pm 2.76$	0.21 $\pm 0.25$
B	40	3.01 $\pm 2.36$	1.42 $\pm 1.40$	8.36 $\pm 5.12$	0.52 $\pm 0.43$
C	40	2.62 $\pm 2.00$	0.98 $\pm 1.04$	5.57 $\pm 5.12$	0.29 $\pm 0.27$
D	40	2.07 $\pm 1.88$	1.07 $\pm 1.10$	8.17 $\pm 7.18$	0.37 $\pm 0.30$
Total	160	2.40 $\pm 2.02$	1.07 $\pm 1.14$	7.15 $\pm 5.96$	0.36 $\pm 0.33$

\*Zn: zinc; Cu: copper; Pb: lead; Cd: cadmium

Table 2- Descriptive data of the lactating women and their infants

	Result	Range
Mothers age (years)	26.4 $\pm$ 4.8	18-41
Infant age (month)	5.6 $\pm$ 5.2	0-22
Height at birth (cm)	50.5 $\pm$ 2.96	42-69
Weight at birth (kg)	3.26 $\pm$ 0.45	2.45-5.25
Head circumference at birth (cm)	34.58 $\pm$ 2.43	23.20-38

In our study, zinc concentration (2.40  $\pm$  2.02 mg/l) was comparable to 4.1 mg/l that was reported in breastmilk of 12-20 weeks postpartum mothers in Austria [14], 1.17–5.31 mg/l in breastmilk of Canadian mothers [15], and 2.0 mg/l in breastmilk of women in Texas [16]. Copper concentration of our study (1.07  $\pm$  1.14 mg/l) was higher than the amounts reported as 0.09–0.27 mg/l in Taiwan [17], 0.36–0.47 mg/l in Korea [18], and 0.86 mg/l in Austria [14]. Our result of lead concentration (7.15  $\pm$  5.96 mg/l) was higher than 0.9  $\pm$  0.4 µg/l in Sweden [19] and 0.55 µg/l in Australia, and lower than 20  $\pm$  5 mg/l in Greece [20], and 35.8  $\pm$  15.0 mg/l in Austria [21]. For cadmium, our results (0.36  $\pm$  0.33 µg/l) were lower than Saudi Arabia (1.73 µg/l) [22], similar

to those found in Japan ( $0.27 \pm 1.82 \mu\text{g/l}$ ) [9] and Slovakia ( $0.43 \pm 0.27 \mu\text{g/l}$ ) [23], and significantly lower than concentrations found in England ( $17.3 \pm 4.9 \mu\text{g/l}$ ) and Germany ( $24.6 \pm 7.3 \mu\text{g/l}$ ). The differences found among various studies might be due to the different assays employed other than variable concentrations of the heavy metals absorbed in body. Entrance of cadmium and lead to breastmilk is universal problem. These harmful heavy metals are abundantly present in the environment and people are exposed to them inevitably [24]. Chou et al. reported that infants of smoking mothers are at higher risk of exposure to cadmium than the infants of nonsmokers [25].

Comparison of our results to other studies is presented at Table 3. We found significant correlation of breastmilk zinc, copper, lead and cadmium with height at birth, weight at birth, head circumference at birth and growth of infants. There was direct correlation between height at birth with zinc concentration but indirect correlation with copper, lead and cadmium in breastmilk. Orun et al. reported that breastmilk cadmium in second month after birth was negatively correlated with z-scores of head circumference and weight of girls at birth while no correlation was observed between lead/cadmium and Edinburgh Postpartum Depression Scale scores [26].

Table 3- Heavy metals concentration in breastmilk of women from different studies

Country	Zn (mg/l)	Cu (mg/l)	Pb (mg/l)	Cd (mg/l)	References
Iran	$2.40 \pm 2.02$	$1.07 \pm 1.14$	$7.15 \pm 5.96$	$0.36 \pm 0.33$	Current study
Iran			3.06–19.47	0.45–5.87	[33]
Iran	2.95	0.36			[34]
Slovakia			4.7	0.43	[35]
Sweden	1.24–5.71	0.32–0.67	0.74–6.4	0.028–0.27	[36]
Spain			0.1–32.3	0.6–11.3	[37]
Portugal	0.39–5.09	0.33–0.97	0.07–4.03		[38]
Croatia	0.62–15.0	0.27–1.35			[39]
Greece	4.90	0.38	0.48	0.19	[3]
Nigeria	0.7	0.83	8.7	9.7	[40]
Kuwait	1.7–3.2	0.41–0.71			[41]
China	40.6	0.67			[42]

De Felip et al. observed no significant association of seafood consumption by Italian mothers with the environmental pollutants in their breastmilk. In addition, no health risk in breastfed infants was observed [27].

Tanner et al. reported that selenium correlates with copper positively [28]. In accordance, Kantola et al. showed that selenium addition to fertilizers in Finland increased selenium concentration in human milk and possibly had impact on zinc and copper concentration in maternal breastmilk [29]. A comparable study was done on lactating cows exposed to environmental toxicants through which increased lead and cadmium were observed in their blood and milk [30].

In one study conducted in Tehran in 2020, amou-

nts of trace elements in raw and bottled milks was measured by DPASV. Concentration of zinc, copper, lead, and cadmium were estimated as  $4.990 \pm 6.244$ ,  $2.424 \pm 4.017$ ,  $0.271 \pm 0.640$  and  $0.099 \pm 0 \mu\text{g/ml}$  in raw milk, respectively, and  $0.999 \pm 0.873$ ,  $0.228 \pm 0.188$ ,  $0.048 \pm 0.033$  and  $0.049 \pm 0 \mu\text{g/ml}$  in bottled milk, respectively [31]. In 2014, heavy metals were measured by DPASV in four brands of baby food (containing rice and wheat) and powdered milk. Concentration of zinc, copper, lead, and cadmium in baby foods ( $n = 240$ ) were  $11.86 \pm 1.474 \text{ mg/100 g}$ ,  $508.197 \pm 83.154 \mu\text{g/100 g}$ ,  $0.445 \pm 0.006 \text{ mg/kg}$ , and  $0.050 \pm 0.005 \text{ mg/kg}$ , respectively. Moreover, their concentration was  $3.621 \pm 0.529 \text{ mg/100 g}$ ,  $403.822 \pm 133.953 \mu\text{g/100 g}$ ,  $0.007 \pm 0.003 \text{ mg/kg}$ ,

and  $0.060 \pm 0.040$  mg/kg, respectively, in powdered milk ( $n = 240$ ) [32].

#### 4. Conclusion

Our investigation showed that breastmilk of Iranian mothers had enough essential elements of zinc and copper. Fortunately, cadmium concentration in the samples was within the safety limits recommended by international agencies. In comparison, lead concentration of breastmilk was high that is a major hazard in public health especially for neonatal and children undergrown at industrial regions. We conclude that there is a need for conducting risk assessment studies to establish national safe limits for heavy metals in breastmilk. Although, more extensive studies with larger sample size should be done to minimize the uncertainties.

#### 5. Acknowledgment

This study was supported by a grant (no. 89-03-33-11269) from Tehran University of Medical Science, Tehran, Iran.

#### 6. Conflict of interest

The authors declare that there is no conflict of interest.

#### References

1. Dorea JG. Zinc in human milk. *Nutrition Research*. 2000; 20(11): 1645-1687.  
[https://doi.org/10.1016/S0271-5317\(00\)00243-8](https://doi.org/10.1016/S0271-5317(00)00243-8)
2. Mead MN. Contaminants in human milk: weighing the risks against the benefits of breastfeeding. *Environmental Health Perspectives*. 2008; 116(10): 427-434.
3. Leotsinidis M, Alexopoulos A, Kostopoulou-Farri E. Toxic and essential trace elements in human milk from Greek lactating women: Association with dietary habits and other factors. *Chemosphere*. 2005; 61: 238-247.  
<https://doi.org/10.1016/j.chemosphere.2005.01.084>
4. Mahan LK, Raymond JL. *Krause's food & the nutrition care process*. 14<sup>th</sup> edition. Elsevier. 2017.
5. Domellof M, Lonnerdal B, Dewey KG. Iron, zinc, and copper concentrations in breast milk are independent of maternal mineral status. *The American Journal of Clinical Nutrition*. 2004; 79: 111-115.

<https://doi.org/10.1093/ajcn/79.1.111>

6. Chen A, Dietrich KN, Huo X, Ho SM. Developmental neurotoxicants in e-waste: an emerging health concern. *Environmental Health Perspectives*. 2011; 119(4): 431-438.  
<https://doi.org/10.1289/ehp.1002452>
7. National Resources Defense Council. *Healthy milk, healthy baby: chemical pollution and mother's milk*. New York. 2001. Available at: <https://www.nrdc.org/media/2001/010522>
8. Kippler M, Bakhtiar Hossain M, Lindh C, Moore SE, Kabir I, Vahter M, Broberg K. Early life low-level cadmium exposure is positively associated with increased oxidative stress. *Environmental Research*. 2012; 112: 164-170.  
<https://doi.org/10.1016/j.envres.2011.11.012>
9. Honda R, Tawara K, Nishijo M, Nakagawa H, Tanebe K, Saito S. Cadmium exposure and trace elements in human breast milk. *Toxicology*. 2003; 186(3): 255-259.  
[https://doi.org/10.1016/s0300-483x\(03\)00002-7](https://doi.org/10.1016/s0300-483x(03)00002-7)
10. Picciano MF. Pregnancy and lactation: physiological adjustments, nutritional requirements and the role of dietary supplements. *The Journal of Nutrition*. 2003; 133: 1997.  
<https://doi.org/10.1093/jn/133.6.1997S>
11. Gadzhieva A. Iron, copper and zinc determination in wine using flame atomic absorption spectroscopy. *Thermo Fisher Scientific, Cambridge, UK*. 2016. 1-3.
12. Chrastny V, Komarek M. Copper determination using ICP-MS with hexapole collision cell. *Chemical Papers*. 2009; 63 (5): 512-519.  
<https://doi.org/10.2478/s11696-009-0057-z>
13. Minnino S, Wang J. Electrochemical methods for food and drink analysis. *Electroanalysis*. 1992; 4: 835.  
<https://doi.org/10.1002/elan.1140040903>
14. Krachler M, Li FS, Rossipal E, Irgolic KJ. Changes in the concentration of trace elements in human milk during lactation. *Journal of Trace Elements in Medicine and Biology*. 1998; 12: 159-176.  
<https://doi.org/10.1016/S0946-672X>
15. Friel JK, Andrews WL, Jackson SE. Elemental composition of human milk from mothers of premature and full-term infants during the first 3 months of lactation. *Biological Trace Element Research*. 1999; 67: 225-247.
16. Hannan MA, Faraji B, Tanguma J. Maternal milk concentration of zinc, iron, selenium, and iodine and

- its relationship to dietary intakes. *Biological Trace Element Research*. 2009; 127:6–15.  
<https://doi.org/10.1007/s12011-008-8221-9>.
17. Lin TH, Jong YJ, Chiang CH, Yang MH. Longitudinal changes in calcium, magnesium, iron, copper and zinc in breast milk of women in Taiwan over a lactation period of one year. *Biological Trace Element Research*. 1998; 62: 31-41.
18. Ko YS, Om AS. Analysis of the concentrations of trace elements in colostrum, transitional and mature milk of Korean women. *The Journal of Trace Elements in Experimental Medicine*. 1998; 11: 407.
19. Hallen P, Jorhem L, Lagerkvist BJ, Oskarsson A. Lead and cadmium levels in human milk and blood. *Science of the Total Environment*. 1995;166: 149-155.  
[https://doi.org/10.1016/0048-9697\(95\)04523-4](https://doi.org/10.1016/0048-9697(95)04523-4)
20. Nashashibi N, Cardamakis E, Bolbos G, Tzinguonis V. Investigation of kinetic of lead during pregnancy and lactation. *Gynecologic and Obstetric Investigation*. 1999; 48 (3): 158-162.  
<https://doi.org/10.1159/000010164>
21. Plockinger B, Dadak C, Meisinger V. Lead, mercury and cadmium in newborn infants and their mothers. *Zeitschrift fur Geburtshilfe und Perinatologie*. 1993; 197 (2): 104-107.
22. Al-Saleh I, Shinwari N, Mashhour A. Heavy metal concentrations in the breast milk of Saudi women. *Biological Trace Element Research*. 2003; 96(1-3): 21-37.  
<https://doi.org/10.1385/BTER:96:1-3:21>
23. Sternowsky HJ, Wessolowski R. Lead and cadmium in breast milk. Higher levels in urban vs. rural mothers during the first 3 months of lactation. *Archives of Toxicology*. 1985; 57 (1): 41-45.  
<https://doi.org/10.1007/BF00286573>
24. Winiarska-Mieczan A. Cadmium, lead, copper and zinc in breast milk in Poland. *Biological Trace Element Research*. 2014; 157: 36-44.  
<https://doi.org/10.1007/s12011-013-9870-x>
25. Chao HH, Guo CH, Huang CB, Chen PC, Li HC, Hsiung DY, Chou YK. Arsenic, cadmium, lead, and aluminum concentrations in human milk at early stages of lactation. *Pediatrics and Neonatology*. 2014; 55: 127-134.  
<https://doi.org/10.1016/j.pedneo.2013.08.005>
26. Orun E, Yalcın SS, Aykut O, Orhan G, Koc Morgil G, Yurdakok K, Uzun R. Breast milk lead and cadmium levels from suburban areas of Ankara. *Science of the Total Environment*. 2011; 409: 2467-2472.  
<https://doi.org/10.1016/j.scitotenv.2011.02.035>
27. Abballe A, Ballard TJ, Dellatte E, Domenico A, Ferri F, Rita Fulgenzi A, Grisanti G, Iacovella N, Ingelido M, Malisch R, Miniero R, Grazia Porpora M, Risica Serena, Ziemacki G, De Felip E. Persistent environmental contaminants in human milk: concentrations and time trends in Italy. *Chemosphere*. 2008; 73: 220-227.  
<https://doi.org/10.1016/j.chemosphere.2007.12.036>
28. Tanner MS, Bhave SA, Kantarjian AH, Pandit AN. Early introduction of copper contaminated animal milk feeds as possible cause of Indian childhood cirrhosis. *The Lancet*. 1983; 322: 992-995.  
[https://doi.org/10.1016/S0140-6736\(83\)90980-7](https://doi.org/10.1016/S0140-6736(83)90980-7)
29. Kantola M, Vartiainen T. Changes in selenium, zinc, copper and cadmium contents in human milk during the time when selenium has been supplemented to fertilizers in Finland. *Journal of Trace Elements in Medicine and Biology*. 2001; 15: 11-17.  
[https://doi.org/10.1016/s0946-672x\(01\)80020-1](https://doi.org/10.1016/s0946-672x(01)80020-1)
30. Patra RC, Swarup D, Kumar P, Nandia D, Naresha R, Ali SL. Milk trace elements in lactating cows environmentally exposed to higher level of lead and cadmium around different industrial units. *Science of the Total Environment*. 2008; 404: 36-43.  
<https://doi.org/10.1016/j.scitotenv.2008.06.010>
31. Sadeghi N, Behzad M, Razavi Sh, Jannat B, Oveisi MR, Hajimahmoodi M. Measurement of zinc, copper, lead, and cadmium in the variety of packaging milk and raw milk in Tehran markets by anodic stripping voltammetry. *Journal of Chemical Health Risks*. 2020; 10(3): 175-183.  
<https://doi.org/10.22034/jchr.2020.1877927.1048>
32. Sadeghi N, Oveisi MR, Jannat B, Hajimahmoodi M, Behfar A, Behzad M, Norouzi N, Oveisi M, Jannat B. Simultaneous measurement of zinc, copper, lead and cadmium in baby weaning food and powder milk by DPASV. *Iranian Journal of Pharmaceutical Research*. 2014; 13(1): 345-349.
33. Goudarzi MA, Parsaei P, Navebpour F, Rahimi E. Determination of mercury, cadmium and lead in human milk in Iran. *Toxicology and Industrial Health*. 2012; 29(9): 1-4.  
<https://doi.org/10.1177/0748233712445047>
34. Khaghani S, Ezzatpanah H, Mazhari N, Givianrad MH, Mirmiranpour H, Sadrabadi FS. Zinc and copper concentrations in human milk and infant formulas. *Iranian Journal of Pediatrics*. 2010; 20: 53-57.

35. Ursinyova M, Massanova V. Cadmium, lead and mercury in human milk from Slovakia. *Food Additives & Contaminants*. 2005; 22: 579-589.

<https://doi.org/10.1080/02652030500135201>

36. Bjorklund KL, Vahter M, Palm B, Grander M, Lignell S, Berglund M. Metals and trace element concentrations in breast milk of first time healthy mothers: a biological monitoring study. *Environmental Health*. 2012; 11: 92-99.

<https://doi.org/10.1186/1476-069X-11-92>

37. Rodriguez EM, Delgado Uretra E, Diaz Romero C. Concentrations of cadmium and lead in different types of milk. *Z Lebensm Unters Forsch A*. 1999; 208: 162-168.

<https://doi.org/10.1007/s002170050395>

38. Almeida AA, Lopes MPVC, Silva MSA, Barrado E. Trace elements in human milk; correlation with blood levels, inter-element correlations and concentration during the first month of lactation. *Journal of Trace Elements in Medicine and Biology*. 2008; 22: 196-205.

<https://doi.org/10.1016/j.jtemb.2008.03.007>

39. Mandic Z, Mandic ML, Grgic J, Grgic Z, Klapac T, Primorac L, Hasenay D. Copper and zinc content in human milk in Croatia. *European Journal of Epidemiology*. 1993; 13: 185-188.

<https://doi.org/10.2307/3582007>

40. Adesiyan AA, Akiibinu MO, Olisekodiaka MJ, Onuegbu AJ, Adeyeye AD. Concentrations of some biochemical parameters in breast milk of a population of Nigerian nursing mothers using hormonal contraceptives. *Pakistan Journal of Nutrition*. 2011; 10: 249-253.

<https://doi.org/10.3923/pjn.2011.249.253>

41. Al-awadi FM, Srikumar TS. Trace-element status in milk and plasma of Kuwaiti and non-Kuwaiti lactating mothers. *Nutrition*. 2000; 16: 1069-1073.

[https://doi.org/10.1016/S0899-9007\(00\)00426-3](https://doi.org/10.1016/S0899-9007(00)00426-3)

42. Liu KS, Hao JH, Xu YQ, Gu XQ, Shi J, Dai CF, Xu F, Shen R. Breast milk lead and cadmium levels in suburban areas of Nanjing. *Chinese Medical Sciences Journal*. 2013; 28(1): 7-15.

[https://doi.org/10.1016/S1001-9294\(13\)60012-7](https://doi.org/10.1016/S1001-9294(13)60012-7)